

Regional grid refinement: unexpected effects on Gulf Stream path and Atlantic overturning

**Kevin L. Rosa^{1,2*}, Mark R. Petersen¹,
Steven R. Brus³, Darren Engwirda^{4,5}, Kristin E. Hoch¹,
Mathew E. Maltrud³, Luke P. Van Roekel³, Phillip J. Wolfram³,**

¹ Computational Physics and Methods (CCS-2), Los Alamos National Laboratory, Los Alamos, NM, USA

² Graduate School of Oceanography, University of Rhode Island, Narragansett, Rhode Island, USA


³ Fluid Dynamics and Solid Mechanics (T-3), Los Alamos National Laboratory, Los Alamos, NM, USA

⁴ NASA Goddard Institute for Space Studies, New York, NY, USA

⁵ Center for Climate Systems Research, Columbia University, New York, NY, USA


Overview

- Testing new coastal-refined variable resolution mesh
- One goal: Hoping to improve Gulf Stream path and strength



Overview

- Testing new coastal-refined variable resolution mesh
- One goal: Hoping to improve Gulf Stream path and strength



- Showed some improvements (e.g. EKE) but did not fix Gulf Stream bias
- This motivated development of a *new* coastal-refined mesh which shows promising preliminary results



Note: A key difference from Kristen's work


>> Here I'm using realistic atmospheric forcing (CORE v2)

Looking for good agreement with observations and with high-res results from Petersen et al. (2019)

Intro/Motivation

1. Why do we want higher resolution models?
2. What's stopping us from running higher resolution models? And what are some solutions?
3. Ways to design an unstructured mesh


1. Why do we want higher resolution models?



Gulf Stream surface relative vorticity - 1.5 km ROMS simulation.

source: Jonathan Gula, Université de Bretagne Occidentale
<http://stockage.univ-brest.fr/~gula/movies.html>

2. What's stopping us from running higher resolution models?



momentum equation:

$$\frac{\partial \mathbf{u}}{\partial t} + \eta \mathbf{k} \times \mathbf{u} + w \frac{\partial \mathbf{u}}{\partial z} = -\frac{1}{\rho_0} \nabla p - \frac{\rho g}{\rho_0} \nabla z^{mid} - \nabla K + \mathbf{D}_h^u + \mathbf{D}_v^u + \mathcal{F}^u$$

thickness equation:

$$\frac{\partial h}{\partial t} + \nabla \cdot (h \bar{\mathbf{u}}^z) + w|_{z=s^{top}} - w|_{z=s^{bot}} = 0$$

tracer equation:

$$\frac{\partial}{\partial t} h \bar{\varphi}^z + \nabla \cdot (h \bar{\varphi} \bar{\mathbf{u}}^z) + \varphi w|_{z=s^{top}} - \varphi w|_{z=s^{bot}} = D_h^\varphi + D_v^\varphi + \mathcal{F}^\varphi$$

hydrostatic condition:

$$p(x, y, z) = p^s(x, y) + \int_z^{z^s} \rho g dz'$$

equation of state:

$$\rho = f_{eos}(\Theta, S, p)$$


Primitive Equations (incompressible hydrostatic Boussinesq)
MPAS-Ocean Model User's Guide 2.0 (2013)

Chessboard graphic:


<http://mathworld.wolfram.com/Chessboard.html>

2. What's stopping us from running higher resolution models?

Low-resolution ocean



4x higher resolution



16x more cells , 4x smaller timestep -> **64x higher computational cost**

>> A low-resolution simulation that runs in a day now takes 2 months to run

2. What's stopping us from running higher resolution models?

Table 1. Setup and performance



	Low-resolution	Coastal-refined	High-resolution
Mesh name	EC60to30	CUSP8	RRS18to6
Horizontal Grid Cells (ocean)	235k	645k	3.69 mil
Cell Size: min-max	30-60 km	8-60 km	6-18 km
Vertical Layers	60	60	80
Time step	30 min	10 min	6 min
Simulated years per day	13.18	4.55	0.77
Total cores (ocean + sea ice + coupler)	960	2160	3600
Million CPU hours per century	0.17 ^a	1.1 ^b	11.9 ^b
Cost vs. low-resolution	×1.0	×6.5	×65.9

^acompy mcnodface

^bblues

2. What's stopping us from running higher resolution models?

**Regional resolution
refinement**




2. What's stopping us from running higher resolution models?

Table 1. Setup and performance


	Low-resolution	Coastal-refined	High-resolution
Mesh name	EC60to30	CUSP8	RRS18to6
Horizontal Grid Cells (ocean)	235k	645k	3.69 mil
Cell Size: min-max	30-60 km	8-60 km	6-18 km
Vertical Layers	60	60	80
Time step	30 min	10 min	6 min
Simulated years per day	13.18	4.55	0.77
Total cores (ocean + sea ice + coupler)	960	2160	3600
Million CPU hours per century	0.17 ^a	1.1	11.9 ^b
Cost vs. low-resolution	×1.0	×6.5	×65.9

^acompy mcnodface
^bblues


3. Ways to design an unstructured mesh




MPAS-Ocean standard meshes



Scaled by observed SSH variability
Sein et al. (2016)



Scaled by Rossby Radius
Sein et al. (2017)



60 km global, 15 km Southern Ocean
Rosa et al. (2018) AGU Poster

3. Ways to design an unstructured mesh



Low-resolution



Coastal-refined



High-resolution



Cost:


x1

x7

x66

Results part 1: California Upwelling

Motivation:




Messié and Chavez (2014)


<1% of ocean area supports:

- * 5% of marine primary production (Carr, 2002) and
- * 20% of fisheries catch (Chavez and Messié, 2009)

Results part 1: California Upwelling





Results part 1: California Upwelling



Results part 1: California Upwelling

Wavenumber power spectral analysis: Quantifying what we saw by eye




Results part 2: Western boundary current

Motivations:

- * Path: Low-resolution MPAS-O (and many other climate models) has unrealistic Gulf Stream (GS) path.

>> Large SST bias in western North Atlantic




SST bias: Low-res model - Observations
Petersen et al. (2019)

Florida-Bahamas Transport Petersen et al. (2019)


EC60to30	RRS18to6	Observations
17.6 Sv	30.1 Sv	31.5 Sv

- * Transport: Low-res GS transport is much weaker than high-res and observations.


Results part 2: Western boundary current




Results part 2: Western boundary current




Results part 2: Western boundary current





Results part 2: Western boundary current





Going to build this figure up piece by piece




Results part 2: Western boundary current




Results part 2: Western boundary current




Results part 2: Western boundary current




44
- 28
- 16
—
= 0





1. Deep Western Boundary Current



(Jack Cook, WHOI Graphics Services)





Fig. 6. Circulation pattern in meridionally bounded ocean with concentrated source S_0 at North Pole and a uniformly distributed sink Q_0 , such that $S_0 = Q_0 a^2 (\phi_2 - \phi_1)$.

Stommel and Arons (1959a)




(Eric S. Taylor, WHOI Graphics Services. Base map from NOAA)

2. Wind-driven Gyre

THE WESTWARD INTENSIFICATION OF WIND-DRIVEN OCEAN CURRENTS


Henry Stommel

(Contribution No. 408, Woods Hole Oceanographic Institution)





Stewart (2008) Fig 11.5 recreation of Stommel (1948) Fig. 4+5

https://www.colorado.edu/oclab/sites/default/files/attached-files/stewart_textbook.pdf





Deep Western Boundary Current	Sub-tropical Gyre
Southward transport	Southward transport
Bouyancy-driven	Wind-driven
Narrow, fast current	Broad; slow speeds
Coastal-refined resolution	Coarse resolution



Results part 2: Western boundary current





Results part 2: Western boundary current





Results part 2: Western boundary current



Results part 2: Western boundary current




Results part 2: Western boundary current




Low-resolution


Mixed Layer Depth







Surface Salinity



Coastal-refined




High-resolution




Low-resolution



Mixed Layer Depth






Surface Salinity




Coastal-refined




High-resolution




Gulf Stream path influenced by resolution?



Gulf Stream path influenced by resolution?



Results part 2: Western boundary current





MERCINA Working Group (2012)

Tail of the Grand Banks:
key "pinch point"

GS / Labrador Current interactions: Salinity at 250 m


GOOD



GS / Labrador Current interactions: Salinity at 250 m



GOOD

High-resolution G case



BAD

Coastally-refined G case



REVIEW



Gulf Stream path being steered by mesh transition






GS impinging on Labrador Current

Weak Gulf Stream

Freshening of Lab Sea

Weak DWBC

Low deepwater formation




Ongoing work


Old




New




Extending refinement around Gulf Stream Extension



Ongoing work




Preliminary results: Improved path of Gulf Stream Extension



Ongoing work

Why is the Gulf Stream path being affected by the mesh resolution transition?







8 km resolution.
Low viscosity



60 km resolution.
High viscosity







8 km resolution.
Low viscosity



60 km resolution.
High viscosity







8 km resolution.
Low viscosity



60 km resolution.
High viscosity



Directed North (higher f) to
conserve Potential vorticity



$$\frac{\zeta + f}{H}$$

Thank you

